



## Praseodymium orthovanadate-waste PET bottle derived sulfur doped carbon for efficient Z-scheme photocatalytic CO<sub>2</sub> reduction

K. Yogesh Kumar<sup>a</sup>, L. Parashuram<sup>b</sup>, M.K. Prashanth<sup>c</sup>, H. Shanavaz<sup>a</sup>, Fahd Alharethy<sup>d</sup>, M. S. Raghu<sup>e,\*</sup>, Byong-Hun Jeon<sup>f,\*</sup>

<sup>a</sup> Department of Chemistry, Faculty of Engineering and Technology, Jain University, Bangalore 562112, India

<sup>b</sup> Department of Chemistry, Nitte Meenakshi Institute of Technology, Yelahanka, Bangalore 560064, India

<sup>c</sup> Department of Chemistry, BNM Institute of Technology, Banashankari, Bangalore 560070, India

<sup>d</sup> Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

<sup>e</sup> Department of Chemistry, New Horizon College of Engineering, Outer Ring Road, Bangalore 560103, India

<sup>f</sup> Department of Earth Resources and Environmental Engineering, Hanyang University, 222, Wangsimni-ro, Seongdong-gu, Seoul 04763, the Republic of Korea

### ARTICLE INFO

#### Keywords:

PET bottle

PrVO<sub>4</sub>

S doped Carbon

Photocatalysis

CO<sub>2</sub> reduction

### ABSTRACT

Escalating amounts of carbon dioxide (CO<sub>2</sub>) in the Earth's atmosphere are a major cause of climate change and its negative consequences, making it a global concern. The present work highlights the intersection of waste management, resource utilization, and climate change mitigation, showcasing the potential for an additional sustainable and circular economy. Synthesizing carbon from waste polyethylene terephthalate (PET) bottles is an eco-friendly approach. X-ray diffraction, Raman spectroscopic, scanning electron microscopic, tunnelling electron microscopic, infrared fourier transform spectroscopy (FTIR) and X-ray photoelectron spectroscopic results indicate the effective extraction of carbon doped with sulfur (S@C) and the successive formation of nanocomposite with PrVO<sub>4</sub> (PrV/S@C). Superior photocatalytic activity was observed under visible light compared to UV light. PrV/S@C showed enhanced light-driven CO<sub>2</sub> reduction compared to pristine materials and was found to evolve 39, 56, and 98 μmol h<sup>-1</sup> g<sup>-1</sup> of H<sub>2</sub>, CH<sub>4</sub> and CO, respectively. The effect of reaction temperature was examined and optimized at 100 °C. The enhanced activity in PrV/S@C compared to PrV and S@C probably due to the increased surface area, conductivity and synergy. In-Situ Diffused reflectance (DRIFT-IR) and liquid chromatography mass spectroscopy (LC-MS) confirm the evolution of CO and CH<sub>4</sub>. The optical, photo/electrochemical, Scavenger studies, Electron spin resonance (ESR) and Mott-Schottky studies suggest the formation of a direct Z-scheme photoredox reaction in the presence of PrV/S@C and sacrificial gent. The reusability studies for 5 cycles indicate good stability with a slight deviation in structure and morphology of PrV/S@C, as observed in XRD and SEM. Tackling plastic pollution and generating value-added carbonaceous products through CO<sub>2</sub> reduction is the twin application of the present work that has piqued the interest of several researchers in environmental remediation.

### 1. Introduction

Carbon dioxide (CO<sub>2</sub>) emissions from the increased use of fossil fuels over the past few decades have resulted in an increase in environmental issues. As human activities continue to emit enormous amounts of CO<sub>2</sub> into the atmosphere, there is an urgent need for new and long-term solutions to lessen the environmental impact [1,2]. Rapid increases in atmospheric CO<sub>2</sub> concentrations have unintended repercussions, including global warming, with temperatures rising at a rate (0.18 °C/decade) that is more than double (0.08 °C/decade) over the previous

100 years [3–5]. The United Nations has pledged to achieve "net zero" emissions by 2050, which means removing as much CO<sub>2</sub> from the atmosphere each year as is emitted due to human activity [6,7]. Attaining this aim will necessitate a global collaboration of scientists, policy-makers, and industries. The development of innovative materials on industrial scales to selectively remove CO<sub>2</sub> from gas mixtures allows for a multifaceted strategy for reducing CO<sub>2</sub> emissions [8,9]. CO<sub>2</sub> can be captured in the atmosphere utilizing materials and processes for biological, chemical, and geological technology, reducing our dependency on fossil fuel supply. Many strategies for turning CO<sub>2</sub> into hydrocarbon

\* Corresponding authors.

E-mail addresses: [dr.msraghu@newhorizonindia.edu](mailto:dr.msraghu@newhorizonindia.edu) (M.S. Raghu), [bhjeon@hanyang.ac.kr](mailto:bhjeon@hanyang.ac.kr) (B.-H. Jeon).

<https://doi.org/10.1016/j.jece.2024.113171>

Received 23 January 2024; Received in revised form 18 April 2024; Accepted 25 May 2024

Available online 31 May 2024

2213-3437/© 2024 Elsevier Ltd. All rights reserved, including those for text and data mining, AI training, and similar technologies.